

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Range expansion of *Ixodes ricinus* to higher altitude, and co-infestation of small rodents with *Dermacentor marginatus* in the Northern Apennines, Italy**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/155832> since 2015-11-23T12:48:46Z

*Published version:*

DOI:10.1016/j.ttbdis.2014.07.021

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



## UNIVERSITÀ DEGLI STUDI DI TORINO

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in *Ticks and Tick-borne Diseases*, Volume 5, Issue 6, October 2014, Pages 970–974. DOI: 10.1016/j.ttbdis.2014.07.021.

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), DOI: 10.1016/j.ttbdis.2014.07.021.

**Range expansion of *Ixodes ricinus* to higher altitude, and co-infestation of small rodents with *Dermacentor marginatus* in the Northern Apennines, Italy.**

Martello Elisa<sup>a</sup>, Mannelli Alessandro<sup>a\*</sup>, Ragagli Charlotte<sup>c</sup>, Ambroggi Cecilia<sup>c</sup>, Selmi Marco<sup>b</sup>,  
Leonardo A. Ceballos<sup>a</sup>, Tomassone Laura<sup>a</sup>

<sup>a</sup> Dipartimento di Scienze Veterinarie, University of Torino, Italy

<sup>b</sup> Osservatorio Permanente per Patologie a trasmissione Vettoriale, ASL2 Lucca, Italy

<sup>c</sup> Ufficio Territoriale per la Biodiversità, Corpo Forestale dello Stato, Lucca, Italy

\*Corresponding author: Alessandro Mannelli, Dipartimento di Scienze Veterinarie, Università degli Studi di Torino, Via Leonardo da Vinci, 44, 10095 Grugliasco (TO), Italy. Tel.: +390116709186; fax: +390116709196; e-mail address: [alessandro.mannelli@unito.it](mailto:alessandro.mannelli@unito.it)

**Abstract.**

Immature ticks (*Ixodes ricinus* and *Dermacentor marginatus*) were collected from small rodents (*Apodemus* spp. and *Myodes glareolus*), in the Northern Apennines, Italy, at an altitude up to 1650 m above sea level (a.s.l.), from 2009 through 2012. While *D. marginatus* had been found at the same location in studies carried out in 1994, *I. ricinus* was very rare or absent. Prevalence (95% confidence interval) of infestation by *I. ricinus* larvae on *Apodemus* spp. was 54.4% (47.5, 61.2), and it was greater than prevalence of *D. marginatus* larvae on the same hosts (23.3%, 17.8, 29.5). The mean (standard deviation) numbers of *I. ricinus* and *D. marginatus* larvae per individual *Apodemus* spp. were similar: 2.3 (4.1) and 2.1 (9.8), respectively. The monthly infestation pattern of the two tick species on *Apodemus* spp. were different. *I. ricinus* larvae were more frequent in June and September, than in July-August. *I. ricinus* nymphs were generally rare, and were most frequently found in July. The prevalence of *D. marginatus* larvae peaked in July-August, whereas nymphs were mostly active in August-September. Increasing population densities of roe deer (*Capreolus capreolus*), and increasing temperatures, in the last decades, in the Apennine area might have contributed to the observed range expansion of *I. ricinus*.

**Keywords**

*Ixodes ricinus*, *Dermacentor marginatus*, *Apodemus* spp., Apennines, Italy.

## Introduction

In the last decades, the geographic range of *Ixodes ricinus*—the most important vector of tick borne zoonoses (TBZ) in Europe—expanded to previously free areas, including greater latitudes in Northern Europe, as well as greater altitudes in the mountains of Central and Southern Europe (Kirby et al., 2004; Materna et al., 2008; Danielová et al., 2006; Jore et al., 2011; Jaenson et al., 2012; Léger et al., 2012). Immature *I. ricinus* are commonly found on small rodents, which may play important roles in the transmission of agents of TBZ. Rodents can also be used as indicators of the presence of the tick and of transmitted agents at a given location (Mannelli et al., 2012a; Pérez et al., 2012; Mihalca and Sándor, 2013).

*Dermacentor marginatus* was the dominant tick species collected on small rodents (*Apodemus* spp. and *Myodes glareolus*) trapped between 1100 and 1400 m a.s.l., in the Northern Apennines, in 1994. In the same study, *I. ricinus* was only found in one out of 128 examined rodents. Furthermore, no host-seeking *I. ricinus* was found by dragging approximately 50 km distance in the same area (Mannelli et al., 1997). In the present article, we report on the analysis of infestation of small rodents by ixodid ticks at the same Apennine location, from 2009 through 2012. The collection of *I. ricinus* from small rodents allowed us to investigate on an increased altitudinal limit for the tick, and to describe the monthly pattern of activity of immature stages, during Summer, in the newly colonized area.

## Materials and methods

### Study area.

The study was carried out on the Tuscan side of the Tuscan-Emilian Apennine National Park, in the province of Lucca, Italy (44° 12' N, 10° 22' E). Climate is characterised by relatively cold winters, and cool summers. In the period 1996-2010, monthly, mean temperature was 1.9 °C in January (minimum), and 17.7 °C in July and August (maximum) (Regione Toscana, 2013a,

2013b). August 2003 was the warmest month on record, with an average temperature of 21.3 °C. Annual rain precipitations are among the greatest in Italy, and showed much variation (between 1350 and 2150 mm) in the 1996-2010 period. Snow cover normally lasts from December through March-April.

Wild ungulates were reintroduced during the 1960's and 1970's. In 2011, the red deer (*Cervus elaphus*) and mouflon (*Ovis orientalis musimon*) populations were estimated at 9.2 and 12.3 head/100 ha. The roe deer was once rare (Ciucci, 1994), but it underwent a remarkable population increase in the last two decades and, in 2011, population density was estimated at 13.2 head/100 ha (Bongi Paolo, unpublished report). The wild boar (*Sus scrofa*) is abundant, although census data are not available.

#### **Small rodent trapping.**

Small rodent trapping was carried out, from 2009 through 2012, during trapping sessions of two or three days, approximately one month apart, from June to September. In 2009, no trapping was carried out in June, whereas repeated trapping sessions were carried out during the other months. No trapping was carried out in September 2011. During the first three years (2009-2011), trapping was carried out in three 3 x 10 grids (30 traps, 10 m apart), for a total of 2520 trap nights. Sites were chosen, within 2 km from the park premises, based upon convenience of access, and at locations where rodents were previously captured in 1994 (Mannelli et al., 1997). Site A, at 1220 m above the sea level (a.s.l.), was characterized by a mixed wood with common hazels (*Corylus avellana*), Turkey oaks (*Quercus cerris*), wild apples (*Malus sylvestris*), and shrubs (*Spartium junceum*, *Rosa canina*). Site B (1140 m a.s.l.) was dominated by oaks, whereas site C (1185 m a.s.l.) was characterized by mixed wood with a predominance of black alders (*Alnus glutinosa*), and including sycamore maples (*Acer pseudoplatanus*), silver firs (*Abies alba*) and European ashes (*Fraxinus excelsior*). Part of the

material collected in 2009-2011 was used in laboratory analysis for the detection of *Rickettsia slovaca* in *D. marginatus* and rodent tissues, as described in Martello et al. (2013). In 2012, trapping was carried out in high stand beech (*Fagus sylvatica*) wood habitat, with poor undergrowth and thick leaf litter, between 1200 and 1650 m a.s.l., near to the altitudinal limit of the tree vegetation (approximately, 1700 m a.s.l.); traps were set 20 m apart in two parallel lines in three transects, for a total of 764 trap nights. Sherman live traps (230x80x90 mm, Sherman Live Traps Co., Tallahassee, FL) and Ugglan live traps (240x60x90 mm, Grahnb, Sweden) were baited with cereals and apples and provided with cotton to protect against the cold temperature during the night.

Captured rodents were anesthetized with a mixture of medetomidine and ketamine, as described in Amore et al. (2007). They were individually identified, in 2009, by ear punch. Starting in 2010, a microchip (transponder AEG ID162 ISO, AEG, Germany) was subcutaneously injected in the interscapular region. Rodent species or genus, sex, and weight were recorded. Each processed animal was carefully screened for the presence of ticks on the entire body. Any attached tick was removed and stored in 70% ethanol. After examination, anesthetized mice were injected with atipamezole HCl (Antisedan®, Pfizer Animal Health, Rome, Italy) to reverse the effects of medetomidine, and were released at the capture site after they had completely recovered from the anaesthesia. Animal capture sampling procedures were approved by the Bioethics Commission of the University of Turin.

#### **Statistical analysis of tick infestation.**

Ticks were identified by species and stage by using keys from Manilla (1998). The prevalence of infestation and 95% confidence interval (CI), were calculated by tick species and stage, year, month, and host species or genus, by using the FREQ procedure in the SAS System 9.3 (SAS, 2011). Data from recapture of the same rodents during the same trapping session were

excluded from the analysis. Mean numbers of ticks per host, standard deviations (SD), and negative binomial dispersion parameters ( $k$ ) were obtained using intercept-only, generalised linear models (GLM) using PROC GENMOD. Negative binomial error (log link) was used to take into account the aggregated distribution of ticks among hosts (Littell et al., 2006).

## Results

### Small rodent trapping.

Trapping yielded 215 *Apodemus* spp. (76 females, 139 males; 80 individuals in 2009, 52 in 2010, 30 in 2011, 53 in 2012), and 57 *Myodes glareolus* (18 females, 39 males; 35 individuals in 2010, 18 in 2011, 4 in 2012). Mean (SD) weight was 24.7 grams (g) (5.9) for *Apodemus* spp., and 27.3g (5.5) for *M. glareolus*. It was not possible to distinguish between the wood mouse *Apodemus sylvaticus* and the yellow-necked mouse *Apodemus flavicollis*. In fact, the distinction between these two species, on the basis of morphological characters, is considered as unreliable, especially in the Southern parts of their geographic range, including central Italy (Barciova and Macholán, 2009; Bugarski-Stanojević et al., 2013; Jojić et al., 2014) .

### Infestation of small rodents by *I. ricinus*.

A total of 485 larvae and 9 *I. ricinus* nymphs were collected from *Apodemus* spp., whereas 17 larvae and one nymph were collected from *M. glareolus*. Prevalence (95% CI) of infestation by *I. ricinus* larvae was 54.4% (47.5, 61.2) on *Apodemus* spp. and 17.5% (8.7, 29.9) on *M. glareolus*. Prevalence of infestation by nymphs was 3.7% (1.6, 7.2) on *Apodemus* spp. and 1.7% (0.04, 9.4) on *M. glareolus*. The mean (SD) number of *I. ricinus* larvae per individual host was 2.3 (4.1) for *Apodemus* spp. and 0.30 (0.73) for *M. glareolus*; the mean (SD) of nymphs was 0.04 (0.21) for *Apodemus* spp. and 0.02 (0.13) for *M. glareolus*. Subsequent statistical analysis was carried out limited to *Apodemus* spp., due to the small sample size for *M. glareolus*.



Prevalence of infestation by *I. ricinus* larvae on *Apodemus* spp. was 48.8% in 2009, 61.5% in 2010, 56.7% in 2011, 54.7% in 2012. *I. ricinus* nymphs were collected from *Apodemus* spp. in 2009, when prevalence was 7.5%, and in 2012, when one individual out of 53 was infested (prevalence = 1.9%). No *I. ricinus* nymphs were found on *Apodemus* spp. in 2010 and 2011.

Infestation levels (prevalence and mean number of ticks per host) by *I. ricinus* larvae on *Apodemus* spp. peaked in June, followed by a decline in July-August, and by a second, lower peak in September (Table 1). Aggregation of *I. ricinus* larvae on *Apodemus* spp. was maximum in August, as shown by the smallest negative binomial parameter,  $k$ , during this month (Table 1). *I. ricinus* nymphs were most frequently found in July (Table 1).

#### **Infestation of small rodents by *Dermacentor marginatus*.**

A total of 442 *D. marginatus* larvae and 79 nymphs were collected from *Apodemus* spp., 34 larvae and 19 nymphs from *M. glareolus*. Prevalence (95% CI) of infestation by *D. marginatus* larvae was 23.3% (17.8, 29.5) on *Apodemus* spp. and 12.3% (5.1, 23.7) on *M. glareolus*. Prevalence of infestation by nymphs was 21.1% (11.4, 33.9) on *Apodemus* spp. and 13.9% (9.6, 19.3) on *M. glareolus*. The mean (SD) number of *D. marginatus* larvae per individual host was 2.1 (9.8) for *Apodemus* spp. and 0.60 (3.0) for *M. glareolus*, whereas the mean (SD) of nymphs was 0.37 (1.2) for *Apodemus* spp. and 0.33 (0.83) for *M. glareolus*.

Prevalence of infestation by *D. marginatus* larvae on *Apodemus* spp. was greater in 2009 (37.2%) and in 2012 (26.4%) than in 2010 (11.5%) and in 2011 (16.7%). The prevalence of infestation by nymphs *D. marginatus* on *Apodemus* spp. was 17.5% in 2009, 11.5% in 2010, 0.0% in 2011, 18.9% in 2012.

Greatest prevalences of infestation by *D. marginatus* larvae on *Apodemus* spp. were observed in July and August. The mean number of larvae per host peaked in July (Table 2). In June, the negative binomial parameter  $k$  was small, indicating a high degree of aggregation (Table 2).

Lowest infestation levels by *D. marginatus* larvae were observed in September, when the distribution was also aggregated. *D. marginatus* nymphs were not found on *Apodemus* spp. in June, and infestation levels by this stage were greatest in late summer, with a peak in August. Aggregation of nymphs among *Apodemus* spp. was maximum in July, when infestation was generally low (Table 2).

#### **Infestation of small rodents by other tick species.**

Two *Ixodes trianguliceps* larvae were found on one *Apodemus* spp. and one *M. glareolus*, whereas two *I. trianguliceps* nymphs were found on one *M. glareolus*.

#### **Discussion**

The analysis of the infestation of small rodents in the period 2009-2012 demonstrated that *I. ricinus* invaded an area in the Northern Apennines of 1100-1650 m a.s.l., where investigations, which were carried out in 1994 showed that this tick species was rare or absent (Mannelli et al., 1997). Increasing population densities of wild ungulates, particularly roe deer (Carnevali et al., 2009) might have played a central role in the observed altitudinal range expansion of *I. ricinus* in our study area. This is in agreement with Medlock et al. (2013), who underlined the importance of changes in distribution and abundance of tick hosts as key determinants of vector range expansion. In the Apennines, increasing ungulate populations and reforestation were associated with the abandonment of traditional land use practices, such as sheep and dairy cow farming (Ciancio et al., 2006).

Climatic factors might be relevant for the expansion of *I. ricinus* at the limits of its altitudinal and latitudinal ranges (Medlock et al., 2013; Materna et al., 2008). Increasing summer temperatures occurred in the Northern Apennines in the last decades, as part of a general warming in the Mediterranean area (Giorgi, 2006; Baldi et al., 2006; Giorgi and Lionello, 2008),

and this might have favored development and activity of *I. ricinus* at greater altitudes. Furthermore, milder winters, with reduced snow cover, might have favored roe deer populations (Gray et al., 2009). Interactions between land use and climate changes might, therefore, have contributed to the observed altitudinal range expansion of *I. ricinus* at our study area.

The time pattern of infestation of small rodents by immature ticks might be the combined result of the effects of factors associated with both the parasites, and the hosts (Rosà et al., 2007; Paulauskas et al., 2009; Paziewska et al., 2010; Pérez et al., 2012; Estrada-Peña et al., 2013). As a consequence, conclusions on seasonal activity of ticks, based upon rodent infestation, must be drawn with caution. In our study, the monthly activities of larvae of *I. ricinus* and *D. marginatus* on the same *Apodemus* spp. hosts, were different, probably reflecting marked differences in the life cycles of these two tick species. The observed peak of *I. ricinus* larvae on *Apodemus* spp. in June were probably derived from egg batches laid by females that fed in the spring and summer of the previous year. Gray (1982) working in Ireland showed that larvae from such egg batches became active in mid-late May and that the majority remained questing at a low level in the vegetation, possibly an adaptation to parasitizing rodents. In our study, *I. ricinus* larval activity declined in July-August, and it increased again in September. These larvae are presumably derived from autumn-feeding females that hatch from winter-diapausing eggs in late summer of the following year (Gray 1982).

The low levels of nymphal *I. ricinus* on small rodents at our study site, might be due to factors leading to a low attachment rate of this stage to small rodents (Gray et al., 1999; Amore et al., 2007). The collection of *I. ricinus* nymphs from the environment, and from host species other than rodents, would provide useful, and complementary information.

Small rodents were found infested by immature *D. marginatus*, confirming previous observations at the same location (Mannelli et al., 1997). Several ungulate species may serve as hosts for adult *D. marginatus*, wild boars being particularly important in the Northern Apennines (Selmi et al., 2008). The seasonal pattern of *D. marginatus* larvae on *Apodemus* spp. was characterized by greatest prevalence of infestation by larvae in July-August, and by peaks of the numbers of larvae per host in June-July, whereas nymphs were active in August-September (Table 2). These observations were in agreement with the relatively short life cycle of the tick species, whose immature stages are usually active and develop during the warmer month of the same year (Nosek et al., 1967; Cringoli et al., 2005). Winter survival of *D. marginatus* in the Northern Apennines is probably associated with the capability of adults to tolerate low temperatures (Dörr and Gothe, 2001).

Aggregation of immature ticks on rodents has been observed previously (Perkins et al., 2003; Kiffner et al., 2011). Factors associated with aggregation of ticks on a small fraction of hosts include sex, movements, home range and grooming activity, as well as the patchy distribution of host-seeking ticks in the environment (Hughes and Randolph, 2001; Mannelli et al., 2012b). The role of individual factors, affecting infestation levels by ticks of different species on small rodents, deserves further investigations given their potential consequences for the transmission of agents of TBZ (Martello et al., 2013).

Levels of infestation by immature ticks (especially *I. ricinus*) were generally low on *M. glareolus*. This host species' relatively small home range, as well as its apparent ability to develop immunity against *I. ricinus*, have been previously suggested as an explanation of lesser tick infestation on *M. glareolus* compared with *Apodemus* spp. (Dizij and Kurtenbach, 1995; Boyard et al., 2008).

The present study was carried out in a limited study area. Therefore, the altitudinal range expansion of *I. ricinus*, and the monthly pattern of activity of immature stages should be

investigated at other locations in the Northern Apennines. Furthermore, trends and patterns of infestation of small rodents by *I. ricinus* and *D. marginatus*, and potential consequences on the transmission dynamics of agents of TBZ, should be followed in time.

### **Conflict of interest**

No competing financial interests exist.

### **Acknowledgements**

We thank Jeremy Gray, University College Dublin, for help in the interpretation of our results on *I. ricinus* monthly patterns. The Corpo Forestale dello Stato, Ufficio Territoriale per la Biodiversità, Lucca, provided accommodation, animal handling facilities, and valuable help. Michela Adami, Donal Bisanzio, Virginia Filipello, and Maurizio Pieroni, helped with fieldwork. The study was funded by the Fondazione Cassa di Risparmio di Lucca, and by the Department of Veterinary Sciences, University of Torino.

## References

- Amore, G., Tomassone, L., Grego, E., Ragagli, C., Bertolotti, L., Nebbia, P., Rosati, S., Mannelli, A., 2007. *Borrelia lusitaniae* in immature *Ixodes ricinus* (Acari: Ixodidae) feeding on common wall lizards in Tuscany, central Italy. J. Med. Entomol. 44, 303-307.
- Baldi, M., Dalu, G., Maracchi, G., Pasqui, M., Cesarone, F., 2006. Heat waves in the Mediterranean: a local feature or a larger-scale effect? Int. J. Climatol. 26, 1477-1487.
- Barciova, L. and Macholán, M., 2009. Morphometric key for the discrimination of two wood mice species, *Apodemus sylvaticus* and *A. flavicollis*. Acta Zoologica Academiae Scientiarum Hungaricae. 55, 31-38.
- Boyard, C., Vourch, G., Barnouin, J., 2008. The relationships between *Ixodes ricinus* and small mammal species at the woodland-pasture interface. Exp. Appl. Acarol. 44, 61-76.
- Bugarski-Stanojević V., Blagojević J., Adnađević T., Jovanović V., Vujošević M., 2013. Identification of the sibling species *Apodemus sylvaticus* and *Apodemus flavicollis* (Rodentia, Muridae) — Comparison of molecular methods. Zool. Anz. 252, 579-587.
- Carnevali, L., Pedrotti, L., Riga, F., Toso, S., 2009. Ungulates in Italy. Status, distribution, abundance, management and hunting of ungulate populations in Italy. Istituto Superiore per la Protezione e la Ricerca Ambientale (ISPRA), 2001-2005 Report. Rome, (<http://www.isprambiente.gov.it/contentfiles/00004300/4327-banca-dati-ungulati-1.pdf/view>).
- Ciancio, O., Corona, P., Lamonaca, A., Portoghesi, L., Travaglini, D., 2006. Conversion of clearcut beech coppices into high forests with continuous cover: a case study in central Italy. Forest Ecol. Manag. 224, 235-240.
- Ciucci, P., 1994. Movimenti, attività e risorse de lupo (*Canis lupus*) in due aree dell'Appennino centro-settentrionale. PhD thesis. Scuola di dottorato in biologia evoluzionistica. University of Rome, La Sapienza.

- Cringoli, G., Iori, A., Rinaldi, L., Veneziano, V., Genchi, C., 2005. Zecche. Rolando Editore, Napoli, Italy.
- Danielová, V., Rudenko, N., Daniel, M., Holubová, J., Materna, J., Golovchenko, M., Schwarzová, L., 2006. Extension of *Ixodes ricinus* ticks and agents of tick-borne diseases to mountain areas in the Czech Republic. Int. J. Med. Microbiol. 296 (Suppl.), 48-53.
- Dizij, A. and Kurtenbach, K., 1995. *Clethrionomys glareolus*, but not *Apodemus flavicollis*, acquires resistance to *Ixodes ricinus* L., the main European vector of *Borrelia burgdorferi*. Parasite Immunol. 17, 177-183.
- Dörr, B. and Gothe, R., 2001. Cold-hardiness of *Dermacentor marginatus* (Acari: Ixodidae). Exp. Appl. Acarol. 25, 151-169.
- Estrada-Peña, A., Gray, J.S., Kahl, O., Lane, R.S., Nijhof, A.M., 2013. Research on the ecology of ticks and tick-borne pathogens – methodological principles and caveats. Front. Cell. Infect. Microbiol. 3, 29.
- Giorgi, F., 2006. Climate change hot-spots, Geophys. Research Letters 33, L08707, doi:10.1029/2006GL025734.
- Giorgi, F. and Lionello, P., 2008. Climate change projections for the Mediterranean region. Global Planet Change. 63, 90-104.
- Gray, J., 1982. The development and questing activity of *Ixodes ricinus* L. under field conditions in Ireland. B. Entomol. Res. 72, 263-270.
- Gray, J.S., Kirstein, F., Robertson, J.N., Stein, J., Kahl, O., 1999. *Borrelia burgdorferi* sensu lato in *Ixodes ricinus* ticks and rodents in a recreational park in south-western Ireland. Exp. Appl. Acarol. 23, 717-729.
- Gray, J.S., Dautel, H., Estrada-Peña, A., Kahl, O., Lindgren, E., 2009. Effects of climate change on Ticks and Tick-borne Diseases in Europe. Interdiscip. Perspect. Infect. Dis. 2009: 593232.
- Hughes, V.L. and Randolph, S.E., 2001. Testosterone increases the transmission potential of tick-

- borne parasites. *Parasitology*. 123(Pt. 4), 365-371.
- Jaenson, T.G.T., Jaenson, D.G.E., Eisen, L., Petersson, E., Lindgren, E., 2012. Changes in the geographical distribution and abundance of the tick *Ixodes ricinus* during the past 30 years in Sweden. *Parasit. Vectors*. 5, 8.
- Jojić V., Bugarski-Stanojević V., Blagojević J., Vujošević M., 2014. Discrimination of the sibling species *Apodemus flavicollis* and *A. sylvaticus* (Rodentia, Muridae). *Zool. Anz*. 253, 261-269.
- Jore, S., Viljugrein, H., Hofshagen, M., Brun-Hansen, H., Kristoffersen, A.B., Nygård, K., Brun, E., Ottesen, P., Sævik, B.K., Ytrehus, B.R., 2011. Multi-source analysis reveals latitudinal and altitudinal shifts in range of *Ixodes ricinus* at its northern distribution limit. *Parasit. Vectors*. 4, 84.
- Kiffner, C., Vor, T., Hagedorn, P., Niedrig, M., Rühle, F., 2011. Factors affecting patterns of tick parasitism on forest rodents in tick-borne encephalitis risk areas, Germany. *Parasitol. Res*. 108, 323-335.
- Kirby, A.D., Smith, A.A., Benton, T.G., Hudson, P.J., 2004. Rising burden of immature sheep ticks (*Ixodes ricinus*) on red grouse (*Lagopus lagopus scoticus*) chicks in the Scottish uplands. *Med Vet. Entomol*. 18, 67-70.
- Léger, E., Vourch, G., Vial, L., Chevillon, C., McCoy, K.D., 2012. Changing distributions of ticks: causes and consequences. *Exp. Appl. Acarol*. 59, 219-244.
- Littell, R.C., Milliken, G.A., Stroup, W.W. , Wolfinger, R.D., Schabenberger O., 2006. SAS for Mixed Models. 2nd Edn. SAS® Institute, Cary, USA.
- Manilla, G., 1998. Acari, Ixodida (Fauna d'Italia 36). Ed. Calderoni, Bologna, Italy.
- Mannelli, A., Tolari, F., Pedri, P., Stefanelli, S., 1997. Spatial distribution and seasonality of ticks (Acarina: Ixodidae) in a protected area in the northern Apennines. *Parassitologia*, 39: 41-45.
- Mannelli, A., Martello, E., Tomassone, L., Calzolari, M., Casalone, C., De Meneghi, D., Dottori, M., Estrada-Peña, A, Fabbi, M., Ferreri, L., Ferroglio, E., Luini, M., Nicolau Solano, S., Ortega, C.,



- Pautasso, A., Prati, P., Vesco, U., 2012a. Inventory of available data and data sources and proposal for data collection on vector-borne zoonoses in animals. Supporting publications 234. European Food Safety Authority (EFSA), Parma.
- Mannelli, A., Bertolotti, L., Gern, L., Gray, J., 2012b. Ecology of *Borrelia burgdorferi* sensu lato in Europe: transmission dynamics in multi-host systems, influence of molecular processes and effects of climate change. FEMS Microbiol. Rev. 36, 837-861.
- Martello, E., Selmi, M., Ragagli, C., Ambroggi, C., Stella, M.C., Mannelli, A., Tomassone, L., 2013. *Rickettsia slovaca* in immature *Dermacentor marginatus* and tissues from *Apodemus* spp. in the Northern Apennines, Italy. Ticks Tick Borne Dis. 4, 518-521.
- Materna, J., Daniel, M., Metelka, L., Hrcarik, J., 2008. The vertical distribution, density and the development of the tick *Ixodes ricinus* in mountain areas influenced by climate changes (The Krkonose Mts., Czech Republic). Int. J. Med. Microbiol. 298, 25-37.
- Medlock, J.M., Hansford, K.M., Bormane, A., Derdakova, M., Estrada-Peña, A., George, J.-C., Golovljova, I., Jaenson, T.G.T., Jensen, J.-K., Jensen, P.M., Kazimirova, M., Oteo, J.Aa., Papa, A., Pfister, K., Plantard, O., Randolph, S.E., Rizzoli, A., Santos-Silva, M.M., Sprong, H., Vial, L., Hendrickx, G., Zeller, H., Van Bortel, W., 2013. Driving forces for changes in geographical distribution of *Ixodes ricinus* ticks in Europe. Parasit. Vectors. 6, 1.
- Mihalca, A.D. and Sándor, A.D., 2013. The role of rodents in the ecology of *Ixodes ricinus* and associated pathogens in Central and Eastern Europe. Front. Cell. Infect. Microbiol. 3, 56.
- Nosek, J., Lichard, M., Sztankay, M., 1967. The ecology of ticks in the Tribec and Hronský Inovec Mountains. Bull. World Health Organ. 36 (Suppl.), 49-59.
- Paulauskas, A., Radzijeuskaja, J., Rosef, O., Turcinaviciene, J., Ambrasiene, D., 2009. Infestation of mice and voles with *Ixodes ricinus* ticks in Lithuania and Norway. Estonian J. Ecol. 58, 112.
- Paziewska, A., Zwolińska, L., Harris, P.D., Bajer, A., Siński E., 2010. Utilisation of rodent species by larvae and nymphs of hard ticks (Ixodidae) in two habitats in NE Poland. Exp. Appl. Acarol. 50,

79-91.

Pérez, D., Kneubühler, Y., Rais, O., Gern, L., 2012. Seasonality of *Ixodes ricinus* ticks on vegetation and on rodents and *Borrelia burgdorferi* sensu lato genospecies diversity in two Lyme borreliosis-endemic areas in Switzerland. Vector Borne Zoonotic Dis. 12, 633-644.

Perkins, S.E., Cattadori, I.M., Tagliapietra, V., Rizzoli, A.P., Hudson, P.J., 2003. Empirical evidence for key hosts in persistence of a tick-borne disease. Int. J. Parasitol. 33, 909-917.

Regione Toscana, 2013a. Dati meteo. Stazione Meteo ORECCHIELLA.  
<http://agroambiente.info.arsia.toscana.it>.

Regione Toscana, 2013b. Scheda Stazione 097 – ORECCHIELLA.  
[http://agrometeo.arsia.toscana.it/modules.php?op=modload&name=CF\\_MenuBlock&file=Manager&act=P\\_1\\_1:@99&act2=97](http://agrometeo.arsia.toscana.it/modules.php?op=modload&name=CF_MenuBlock&file=Manager&act=P_1_1:@99&act2=97).

Rosà, R., Pugliese, A., Ghosh, M., Perkins, S.E., Rizzoli, A., 2007. Temporal variation of *Ixodes ricinus* intensity on the rodent host *Apodemus flavicollis* in relation to local climate and host dynamics. Vector Borne Zoonotic Dis. 7, 285-295.

SAS, 2011. SAS/STAT 9.3 User's Guide. SAS® Institute, Cary, USA.

Selmi, M., Bertolotti, L., Tomassone, L., Mannelli, A., 2008. *Rickettsia slovaca* in *Dermacentor marginatus* and tick-borne lymphadenopathy, Tuscany, Italy. Emerg. Infect. Dis. 14, 817-820.

Table 1. Infestation of *Apodemus* spp. by immature *Ixodes ricinus* in the Northern Apennines, Italy, from 2009 through 2012.

Month (n)	June (48)			July (46)			August (72)			September (49)		
Tick stage	Prevalence (95% CI)	Mean (SD)	k	Prevalence (95% CI)	Mean (SD)	k	Prevalence (95% CI)	Mean (SD)	k	Prevalence (95% CI)	Mean (SD)	k
Larvae	68.8 (53.7, 81.3)	3.1 (5.4 )	0.60	50.0 (34.9, 65.1 )	1.4 (1.9 )	0.66	40.3 (28.9, 52.5)	1.6 (3.7 )	0.25	65.3 (50.4, 78.3)	3.2 (4.3)	0.61
Nymphs	2.1 (0.05, 11.1 )	0.02 (0.14)	ND	6.5 (1.4, 17.9)	0.09 (0.35 )	0.14	2.8 (0.34, 9.7)	0.03 (0.17)	ND	2.0 (0.05, 10.9 )	0.02 (0.14)	ND

n: number of examined hosts. Prevalence: percent of examined hosts with a least one tick. 95% CI: exact binomial 95% confidence limits of prevalence. Mean: mean number of ticks per examined host. SD: standard deviation. k: negative binomial distribution parameter, small values indicate aggregation of ticks among hosts. ND: not determined, due to only one or two individual infested by ticks.

Table 2. Infestation of *Apodemus* spp. by immature *Dermacentor marginatus* in the Northern Apennines, Italy, from 2009 through 2012.

Month (n)	June (48)			July (46)			August (72)			September (49)		
Tick stage	Prevalence (95% CI)	Mean (SD)	k	Prevalence (95% CI)	Mean (SD)	k	Prevalence (95% CI)	Mean (SD)	k	Prevalence (95% CI)	Mean (SD)	k
Larvae	8.3 (2.3, 20.0)	2.4 (11.6)	0.02	34.8 (21.4, 50.3)	5.0 (17.0)	0.1	37.5 (26.4, 49.7)	1.1 (2.1)	0.3	6.1 (1.3, 16.9)	0.37 (2.3)	0.02
Nymphs	0.00 (0.00, 7.4)	0.00 (0.00)	ND	4.4 (0.53, 14.8)	0.20 (0.93)	0.02	27.8 (17.9, 39.6)	0.74 (1.6)	0.23	16.3 (7.3, 29.7)	0.35 (1.1)	0.14

n: number of examined hosts. Prevalence: percent of examined hosts with a least one tick. 95% CI: exact binomial 95% confidence limits of prevalence. Mean: mean number of ticks per examined host. SD: standard deviation. k: negative binomial distribution parameter, small values indicate aggregation of ticks among hosts. ND: not determined, due to only one or two individual infested by ticks.